

**KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING**



FLO-MO

Liquid Fertiliser

CROP	APPLICATION RATES AND TIMING		WATER L/Ha
	RATE ml/Ha	TIMING	
Brassicac:	150-250	Minimum of two applications - first at 4-6 leaf stage and repeat as necessary at 7 to 10 day intervals.	200
Cucurbits	100-150	Minimum of two applications - first at 2 to 4 leaf stage and repeat as necessary at 7 to 10 day intervals	200
Lucerne:	150-250	For established stands apply every 3rd or 4th cut. New plantings apply as soon as sufficient leaf cover	100-200
Canola:	200-300	Apply at 3 or 5 leaf stage. Where deficiency is severe a repeat application may be necessary.	100-200
Lettuce:	150-250	Apply first application at 2 to 4 leaf stage and repeat at 7 to 10 day intervals as necessary.	200
Tomatoes:	200	Apply when growth at 10to 15cm.	200
Lupins/Cereals	200-300	Apply at 3 to 5 leaf stage. Where deficiency is severe a repeat application may be necessary.	100-200
Soya Beans			
Sunflowers:	200-400	Apply at 4 to 6 leaf stage.	100-200
Legumes, Broadacre Crops and Pastures:	150-250	Apply Spring and Autumn.	100-200
Vegetables		An initial glasshouse application may be made at the 2 leaf stage of 5ml/1000 plants in 5Lts of water	
Seedlings:			

ANALYSIS

	%W/W	g/L
Molybdenum(Mo) as molybdenum phosphate	25.00	250
Phosphorus (P) as water soluble	11.00	110
Colour stabiliser	0.20	2
Surfactant	0.50	5
This product contains		
Solids	61%w/w	
Water	39%w/w	

Manufactured by
RUTEC PTY LTD
23 HARGRAVE STREET, TAMWORTH NSW 2340

1L 5L 20L

GENERAL INSTRUCTIONS

MIXING AND SPRAYING

Shake the container thoroughly before use. Pour the product through the top filter into the tank three quarters filled with water, whilst agitating. Top up with water and continue to agitate until spraying is complete.

COMPATABILITY

Rutec products are widely compatible with other agrochemicals. Contact your distributor for information.

PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT

Do NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed original container, in a well-ventilated area as cool as possible. Protect from frost. Do not store for prolonged periods in direct sunlight.

Triple rinse containers before disposal and add hoses to tank mix or dispose of rinsate in a disposal pit away from desirable plants, their roots and watercourses.

On-site disposal of undiluted chemicals is unacceptable. Destroy empty containers by breaking, crushing or puncturing them. Dispose of the containers at a local authority landfill that does not burn its refuse. If there is no local authority landfill readily available in your area, bury the containers to a depth of 500mm or more at a licenced/approved disposal site. In some states wastes can only be buried at a licenced landfill. Do not burn empty containers or product.

SPILLAGE

Hose small spillage (less than 1 litre of concentrate) away with a large excess of water. Large spillage should not be directed to drains. Large spillage should be absorbed in sawdust or sand and the absorbent material containing the product disposed of in a sealed container at an approved disposal site.

SAFETY DIRECTIONS

Harmful if swallowed. Avoid contact with eyes, skin and clothing. Wash thoroughly with soap and water after handling. Avoid inhaling spray mist.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre Ph: 13 11 26. If in eyes, flush with water for 15 minutes and see a doctor.

EXCLUSION OF WARRANTIES AND LIMITATIONS OF LIABILITY

The purchaser acknowledges that no express or implied warranty, condition, description or representation is given or made by Rutec Pty Ltd (ABN 91 002 846 988) in relation to this product. To the maximum extent permitted by the law, all warranties, terms, conditions and any duty of care which would otherwise be implied or arise by operation of law, statute or use in relation to this product are hereby excluded as against any and all persons, whether purchasers of this product or not.

To the maximum extent permitted by law, Rutec's total liability to any person in connection with this product, whether arising from a breach of any express or implied term, condition or warranty, any duty of care or otherwise, other than any liability which by these terms or otherwise is excluded, shall not exceed the amount paid by the purchaser to the vendor for this product, whether or not such liability results from or involves negligence on the part of Rutec.

WARNING: Excessive Molybdenum can be harmful to stock - do not apply molybdenum fertiliser to more than 25% of a farm annually. Do not exceed a concentration of 0.25litres of Flo Mo in 100 litres of water.

Withholding period for ruminants - 14 days