

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING

APPLICATION RATES AND TIMING

Apples, Pears: 4-6L/ha. Commence application 4-6 weeks after petal fall. Repeat as necessary at 10-14 day intervals. Water rate 500-1000L/ha

Apricots, Cherries, Peaches, Plums, Nectarines: 4-6L/ha. Commence applications at stone hardening. Repeat applications may be made at 10-14 day intervals up to 1 week before harvest. Water rate 500-1000L/ha

Brassicas, Brussel Sprouts, Cabbage, Cauliflower, Calabrese, Broccoli: 4-6L/ha. Commence application at 4-6 true leaf stage. Repeat as necessary at 7-10 day intervals. Water rate 250-750L/ha.

Carrots: 4-6L/ha. When sufficient leaf area to intercept spray at 10-14 day intervals. Water rate 200-500L/ha

Citrus: 4-6L/ha. Commence applications after fruit set and repeat as necessary at 14-21 day intervals. Water rate- at least 1000L/ha

Cotton: 3-6L/ha. 3 applications at 14 day intervals commencing prior to first flower. Water rate 30-100L/ha

Lucerne: 6L/ha. When sufficient leaf area to intercept spray. Water rate 50-250L/ha

Mangos: 3L/ha. Commence applications at fruit swelling and repeat as necessary at 14-21 day intervals. N.B. Do not apply within 7-10 days of any oil application. Water rate 400-1000L/ha

Onions: 6L/ha. When sufficient leaf area to intercept spray. Water rate 200-500L/ha

Pineapple: 6-10L/ha. Apply 4-6 weeks before harvest. Water rate 400-600L/ha

Potatoes: 4-6L/ha. Apply during bulking and repeat as necessary at 7-10 day intervals. Water rate 200-500L/ha

Strawberries: 6L/ha. Commence after flowering and repeat as necessary at 7-10 day intervals. Water rate 500-1000L/ha

Tomatoes: 4-6 L/ha. Commence applications at flowering of second truss. Repeat as necessary at 7-10 day intervals. Water rate 200-500L/ha

Turf: 20L/ha. As soon as growth commences in Spring. In the case of moderate to severe deficiency repeat applications at 14 day intervals. Water rate 500-1000L/ha

Vines: 4-6L/ha. At fruit set, pea sized berries, and the start of ripening. Water rate 500-1000L/ha

Fertigation: Up to 60 L/ha. Repeat as necessary at 7-14 day intervals



K MIX 320

ANALYSIS

Potassium (K) as potassium citrate

%W/V

g/L

32.00

320

As EDTA complex

Magnesium (Mg)

0.03

0.30

Copper (Cu)

0.016

0.16

Iron (Fe)

0.16

1.60

Manganese (Mn)

0.031

0.31

Zinc (Zn)

0.031

0.31

Boron (B) as borax

0.02

0.20

Molybdenum (Mo) as sodium molybdate

0.016

0.16

Wetting Agent

0.05%w/v

This product contains: Water

16.0%w/w

Solids

84.0%w/w

20L 200L 1000L

Manufactured by

RUTEC PTY LTD

23 HARGRAVE STREET
TAMWORTH NSW 2340

GENERAL INSTRUCTIONS

MIXING AND SPRAYING

Fill the tank to a minimum of half volume. Add compatible chemicals as required maintaining agitation. Fill tank with water while maintaining agitation. Mix enough Rutec K Mix 320 for 1 day's application. Do not store mixed with water or other chemicals.

PRECAUTIONS

Avoid application under extremes of climate eg. Rapid drying conditions, extremely slow drying conditions, frost rain or when frost or rain are anticipated

COMPATIBILITY

RUTEC K Mix 320 is widely compatible with many agrichemicals. When mixing with other chemicals, always mix a small quantity and observe precipitation in mixture. If precipitation occurs DO NOT proceed with application.

STORAGE AND DISPOSAL

Store in the closed original container, in a well-ventilated area as cool as possible. Protect from frost. Do not store for prolonged periods in direct sunlight.

Triple rinse containers before disposal and add hoses to tank mix or dispose of rinsate in a disposal pit away from desirable plants, their roots and watercourses.

On-site disposal of undiluted chemicals is unacceptable. Destroy empty containers by breaking, crushing or puncturing them.

Dispose of the containers at a local authority landfill that does not burn its refuse. If there is no local authority landfill readily available in your area, bury the containers to a depth of 500mm or more at a licensed/approved disposal site. In some states wastes can only be buried at a licensed landfill. Do not burn empty containers or product.

SPILLAGE

Hose small spillage (less than 1 litre of concentrate) away with a large excess of water. Large spillage should not be directed to drains. Large spillage should be absorbed in sawdust or sand and the absorbent material containing the product disposed of in a sealed container at an approved disposal site.

PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT

Do NOT contaminate streams, rivers or waterways with the chemical or used containers.

SAFETY DIRECTIONS

Harmful if swallowed. Avoid contact with eyes, skin and clothing. Wash thoroughly with soap and water after handling. Avoid inhaling spray mist.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre Ph: 13 11 26. If in eyes, flush with water for 15 minutes and see a doctor.

EXCLUSION OF WARRANTIES AND LIMITATIONS OF LIABILITY

The purchaser acknowledges that no express or implied warranty, condition, description or representation is given or made by Rutec Pty Ltd (ABN 91 002 846 988) in relation to this product. To the maximum extent permitted by the law, all warranties, terms, conditions and any duty of care which would otherwise be implied or arise by operation of law, statute or use in relation to this product are hereby excluded as against any and all persons, whether purchasers of this product or not. To the maximum extent permitted by law, Rutec's total liability to any person in connection with this product, whether arising from a breach of any express or implied term, condition or warranty, any duty of care or otherwise, other than any liability which by these terms or otherwise is excluded, shall not exceed the amount paid by the purchaser to the vendor for this product, whether or not such liability results from or involves negligence on the part of Rutec.